



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/83, 15/84, 15/63, 5/10, A01H 4/00		A1	(11) International Publication Number: WO 99/25855 (43) International Publication Date: 27 May 1999 (27.05.99)
(21) International Application Number: PCT/US98/24486 (22) International Filing Date: 16 November 1998 (16.11.98) (30) Priority Data: 60/065,627 18 November 1997 (18.11.97) US 60/065,613 18 November 1997 (18.11.97) US 60/099,461 8 September 1998 (08.09.98) US (71) Applicant (for all designated States except US): PIONEER HI-BRED INTERNATIONAL, INC. [US/US]; 800 Capital Square, 400 Locust Street, Des Moines, IA 50319 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): BASZCZYNSKI, Christopher, L. [US/US]; 7305 Benton Drive, Urbandale, IA 50322 (US). LYZNIK, Leszek, Alexander [PL/US]; 5960 Greendale Place #105, Johnston, IA 50131 (US). GORDON-KAMM, William, J. [US/US]; 3916 67th Street, Urbandale, IA 50322 (US). GUAN, Xueni [CN/US]; 3668 Ruelle de Ville, San Diego, CA 82139 (US). (74) Agents: SPRUILL, W., Murray et al.; Alston & Bird LLP, Bell Seltzer Intellectual Property Law Group, P.O. Drawer 34009, Charlotte, NC 28234 (US).		(81) Designated States: AL, AM, AT, AT (Utility model), AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, CZ (Utility model), DE, DE (Utility model), DK, DK (Utility model), EE, EE (Utility model), ES, FI, FI (Utility model), GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (Utility model), SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(54) Title: MOBILIZATION OF VIRAL GENOMES FROM T-DNA USING SITE-SPECIFIC RECOMBINATION SYSTEMS			
(57) Abstract The invention relates to methods and compositions for site-specific recombinase-mediated mobilization of viral replicons and associated DNAs of interest from T-DNA. The methods of the invention comprise <i>Agrobacterium</i> -mediated transfer recombinase. The nucleic acids of the invention may additionally contain expression cassettes encoding the cognate site-specific recombinase for the target sites flanking the viral genome. The compositions of the invention further comprise <i>Agrobacterium</i> containing the nucleic acids of the invention. The compositions and methods of the invention have use in increasing the efficiency of agroinfection, providing high copy numbers of a DNA of interest for transient expression or for integration into a plant chromosome, and in simplifying the construction and stable maintenance of vectors for agroinfection and transformation.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

-1-

MOBILIZATION OF VIRAL GENOMES FROM T-DNA USING SITE-SPECIFIC RECOMBINATION SYSTEMS

FIELD OF THE INVENTION

The present invention relates to plant molecular biology. More specifically, the invention relates to methods and compositions for increasing the efficiency of excision of viral replicon from T-DNA that is transferred to a plant by agroinfection.

BACKGROUND OF THE INVENTION

Agrobacterium harboring a Ti or a Ri plasmid can efficiently transfer a portion of these plasmids, the T-DNA, into plant cells at the site of a wound. Transfer of the T-DNA into the plant cell is induced by signal compounds present at the site of a plant wound and requires T-DNA border sequences at both ends of the T-DNA and trans-acting virulence gene products (*vir*) encoded by the Ti or Ri plasmid. The transferred T-DNA is then targeted to the nucleus and integrated into the plant genome. A tumor, crown gall, forms at the site of inoculation in most dicotyledonous plants.

Tumor formation results from the expression of T-DNA oncogenes encoding the growth factors auxin and cytokinin which promote plant cell proliferation. In order for these oncogenes to be expressed, the T-DNA must first integrate into the genome of the plant. Tumor formation is limited to dicotyledonous plants because the T-DNA, while transferred into the cells of monocots, is either not normally integrated into the genome of monocotyledonous plants or is integrated and silenced.

By inserting a viral genome into the T-DNA, *Agrobacterium* can be used to mediate the viral infection of plants. Following transfer of the T-DNA to the plant cell, excision of the viral genome from the T-DNA (mobilization) is required for successful viral infection. This *Agrobacterium*-mediated method for introducing a virus into a plant host is known as agroinfection (for a review, see

Grimsley, "Agroinfection" pp. 325-342, in *Methods in Molecular Biology*, vol 44: *Agrobacterium Protocols*, ed. Gartland and Davey, Humana Press, Inc., Totowa, NJ; and Grimsley (1990) *Physiol. Plant.* 79:147-153, the contents of which are incorporated herein by reference). Upon entry into the plant cell nucleus, a unit
5 length circular viral genome that is capable of initiating systemic infection is mobilized from the T-DNA. Integration of the T-DNA into the plant genome is not required for this event. Two non-exclusive mechanisms, intramolecular homologous recombination and replicative release, have been proposed for this release of circular viral genomes or replication intermediates from T-DNA. The
10 replicative release of viral genomes by the rolling-circle replication mechanism has been demonstrated for the mobilization of geminivirus genomes from T-DNA (Stenger *et al.* (1991) *Proc. Natl. Acad. Sci.* 88:8029-8033). Release of geminivirus via homologous recombination between tandemly repeated genomes has also been demonstrated (Lazarowitz *et al.* (1989) *EMBO J.* 8:1023-1032).
15 Release by either of the above mechanisms requires the presence of tandem directly repeated copies of the viral replicon in the T-DNA. A circular viral replicon may be excised from the T-DNA by intramolecular homologous recombination between the repeated genomes. For replicative release, two origin of replication sequences must be present to initiate and complete the replication
20 process. Both mechanisms of mobilization are complex biochemical processes that may be attenuated by a number of factors which in turn affect the efficiency of viral excision. Furthermore, tandem dimers of viral DNA are often difficult to construct and are unstable in recombination proficient host cells.

Agroinfection has been reported in a number of publications as a
25 successful method for inducing systemic viral infections in plant cells, including monocotyledonous plants such as maize (Heath *et al.* (1997) *Mol. Plant-Microbe Interact.* 10:221-227, Grimsley *et al.* (1989) *Mol. Gen. Genet.* 217:309-316, the contents of which are incorporated herein by reference). In many instances, particularly when naked viral nucleic acid is non-infectious, agroinfection is the
30 only way of transforming a plant with cloned viral DNA. Even where naked viral nucleic acid is infectious, agroinfection is frequently used because it is relatively

-3-

efficient and does not require the production of large amounts of plasmid or viral DNA. Agroinfection has been used to study viral replication and recombination, in the investigation of viral gene functions, for the production of autonomously replicating viral vectors, for transient expression of genes inserted into T-DNA, for
5 integration of DNA into a plant genome, for production of virus resistant plants, for the study of transposable elements and for the determination of tissue-specific susceptibility to T-DNA transfer.

The development of plant virus gene vectors for expression of foreign genes in plants provides a means to provide high levels of gene expression
10 within a short time. The benefits of virus-based transient RNA and DNA replicons include rapid and convenient engineering coupled with flexibility for expeditious application in various plant species. In this manner, autonomously replicating viruses offer numerous advantages for use as vehicles for transient expression of foreign genes, including their characteristic high levels of multiplication and
15 concomitant levels of transient gene expression. Accordingly, it would be beneficial to provide methods that facilitate the construction of vectors for agroinfection, provide flexibility in designing viral vectors for genetic transformation of plant cells and increase the efficiency of mobilization of viral replicons from T-DNA and the copy number of a DNA sequence of interest
20 associated with the viral replicon.

The present invention accomplishes these objectives by providing methods and compositions for the mobilization of viral replicon from T-DNA via site-specific recombination systems. Site-specific recombination-mediated excision of DNA fragments from chromosomal or extrachromosomal DNA
25 molecules has been described for a number of site-specific recombination systems and plant species. See Russell *et al.* (1992) *Mol. Gen. Genet.* 234:49-59; Lyznik *et al.* (1996) *Nucleic Acids Res.* 24:3784-3789; and Dale *et al.* (1991) *Proc. Natl. Acad. Sci.* 88:10558-10562 the contents of which are incorporated by reference. However, mobilization of viral vectors from T-DNA via site-specific
30 recombination has not previously been applied to agro-mediated transformation.

SUMMARY OF THE INVENTION

The invention relates to methods and compositions for site-specific recombinase-mediated mobilization of viral replicons and associated DNAs of interest from T-DNA. The methods of the invention comprise *Agrobacterium*-mediated transfer of T-DNA to a plant cell, wherein the T-DNA contains a viral replicon flanked by directly repeated target sites for a site-specific recombinase and optionally a DNA of interest linked to the viral replicon. The DNA of interest may also contain a non-identical target site for the recombinase. An expression cassette for the site-specific recombinase is present on the T-DNA or the plant genome, or is transiently introduced into the plant cell. Expression of the site-specific recombinase in the plant cell results in excision of the viral replicon and the associated DNA of interest. The viral replicon and DNA of interest are then replicated to high copy number in the host plant cell.

The compositions of the invention comprise nucleic acids, such as T-DNAs containing a viral DNA flanked by directly repeated target sites for a site-specific recombinase. The nucleic acids of the invention may additionally contain expression cassettes encoding the cognate site-specific recombinase for the target sites flanking the viral genome. The compositions of the invention further comprise *Agrobacterium* containing the nucleic acids of the invention.

The compositions and methods of the invention have use in providing high copy numbers of a DNA of interest for transient expression or for integration into a plant chromosome, in simplifying the construction and stable maintenance of vectors for agro-mediated transformation of plants and in increasing the efficiency of agroinfection.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 schematically illustrates a model of mobilization of a viral replicon from T-DNA by *loxP*/Cre site-specific recombination.

Figure 2 schematically illustrates Cre-mediated site-specific excision of a viral DNA and DNA of interest from a T-DNA.

Figure 3 schematically illustrates the integration of DNA into a chromosome by double site-specific recombination.

Figure 4 shows the 5' to 3' sequences of the *loxP* and FRT sites.

Figure 5 schematically illustrates a PCR protocol for construction of
5 a viral DNA flanked by *loxP* target sites and XhoI restriction sites.

DETAILED DESCRIPTION OF THE INVENTION

The invention is directed to methods and compositions for the mobilization of viral DNA from T-DNA. In the methods of the invention, a viral
10 replicon flanked by directly repeated target sites for a site-specific recombinase is inserted into a T-DNA carried by an *Agrobacterium* replicon. The T-DNA is transferred into the cell of a plant by agro-mediated transfer. Expression of the site-specific recombinase in the infected plant cell will result in excision of a circular viral replicon from the transferred T-DNA (Figure 1). Replication of this
15 viral replicon will result in a high number of copies of the replicon. Systemic infection of the plant may take place if the replicon encodes infectious viral particles. Accordingly, the methods of the invention have use in the production of viral DNA and/or in promoting systemic viral infection of a plant.

Thus, the invention is directed to a method for mobilizing a viral
20 replicon from a T-DNA, comprising:

- a) providing an *Agrobacterium* replicon having a T-DNA, wherein said T-DNA contains a viral replicon flanked by directly repeated target sites for a site-specific recombinase; and
- b) infecting a cell of a plant with an *Agrobacterium*
25 carrying said *Agrobacterium* replicon under conditions that allow transfer of said T-DNA and expression of said recombinase in said cell;

wherein said cell, said T-DNA, or said viral replicon contains a nucleotide sequence encoding said recombinase or an active fragment or variant thereof, and said nucleotide sequence is operably linked to a promoter that
30 drives expression in said cell.

If a DNA of interest has been inserted into the viral replicon, or

-6-

between the viral replicon and a target site for a site-specific recombinase, the DNA of interest will also be replicated to high copy. The high copy number of the DNA of interest increases the efficiency of integration of this DNA into the genome, or the level of transient expression of a gene encoded by the DNA of interest. Thus, in another aspect, the invention provides a method for providing to
5 a plant cell a plurality of copies of a DNA sequence of interest, comprising:

a) providing an *Agrobacterium* replicon having a T-DNA, wherein said T-DNA contains in a 5' to 3' or 3' to 5' orientation, a first target site for a site-specific recombinase, a viral replicon, said DNA sequence of interest,
10 and a second target site for said recombinase in direct repeat with said first target site, wherein said first and second target sites are identical; and

b) infecting a cell of a plant with an *Agrobacterium* carrying said *Agrobacterium* replicon under conditions that allow transfer of said T-DNA and expression of said recombinase in said cell; wherein said cell, said T-DNA, or said viral replicon contains a nucleotide sequence encoding said
15 recombinase or an active fragment or variant thereof, and said nucleotide sequence is operably linked to a promoter that drives expression in said cell.

The methods of the invention are useful for providing a high copy number of a DNA of interest for site-specific integration into a plant chromosome. In this case, the plant chromosome will contain one or more target sites for a site-specific recombinase. Depending on the substrates, the recombination event could consist of a single or a double cross-over event. In the simplest case, the plant chromosome and the excised viral replicon each contain a single target site. Site-specific recombination between these two target sites results in the insertion of the
20 viral replicon and any associated DNA of interest into the plant chromosome.

The inclusion of a non-identical target site between the viral replicon and the DNA of interest allows integration of the DNA of interest into a plant genome having a corresponding target site, without concomitant insertion of the viral replicon. In this method, the distal ends of the viral replicon and the DNA
30 of interest are flanked by identical target sites. Recombination between the identical target sites results in excision of a circular viral replicon containing the

DNA of interest flanked by non-identical target sites for the recombinase (Figure 2). For targeted insertion of the DNA of interest, the same two non-identical target sites are present in the target organisms genome, thereby establishing a target site for insertion of the DNA of interest. A double crossover event resulting from site-specific recombination between the identical target sites on the host genome and the DNA of interest results in the insertion of the DNA of interest into the chromosome of the target organism, free of the viral replicon (Figure 3). Such methods are discussed in detail in the co-pending provisional application, Serial No 60/065,613, filed 18 November 1997, the contents of which are incorporated herein by reference.

Thus, in a further aspect, the invention provides a method for providing to a plant cell a plurality of copies of a DNA sequence of interest flanked by nonidentical target sites for a site-specific recombinase, comprising:

a) providing an *Agrobacterium* replicon having a T-DNA, wherein said T-DNA contains in a 5' to 3' or 3' to 5' orientation, a first target site for said recombinase, a viral replicon, a second target site for said recombinase, said DNA sequence of interest, and a third target site for said recombinase, wherein said first and third target sites are directly repeated and identical with respect to each other, and said second target site is nonidentical to said first and third target sites; and

b) infecting a cell of a plant with an *Agrobacterium* carrying said *Agrobacterium* replicon under conditions that allow transfer of said T-DNA and expression of said recombinase in said cell;

wherein said cell, said T-DNA, or said viral replicon contains a nucleotide sequence encoding said or an active fragment or variant thereof, and said nucleotide sequence is operably linked to a promoter that drives expression in said cell.

The compositions of the invention comprise T-DNAs containing a viral replicon flanked by directly repeated target sites for a site-specific recombinase. In another aspect, the compositions of the invention comprise a T-DNA containing in a 5' to 3' or 3' to 5' orientation, a first target site for said

recombinase, a viral replicon, a second target site for said recombinase, said DNA sequence of interest, and a third target site for said recombinase, wherein said first and third target sites are directly repeated and identical with respect to each other, and said second target site is nonidentical to said first and third target sites. The T-DNAs of the invention may further comprise nucleotide sequence encoding a site-specific recombinase, wherein the nucleotide sequence is operably linked to a promoter that drives expression in a plant cell. The compositions of the invention further comprise replicons containing these T-DNAs and *Agrobacterium* containing these replicons.

By "mobilization of a viral replicon" is meant excision of a viral replicon from a T-DNA sequence after it has been transferred to a plant cell. In the methods of the invention, mobilization of viral DNA is accomplished by conservative site-specific recombination between directly repeated target sites flanking the viral DNA. In one embodiment, the resulting product is a circularized viral DNA containing one copy of the target site (see Figure 1).

By "viral replicon" is meant double-stranded DNA from a virus having a double stranded DNA genome or replication intermediate. The excised viral DNA is capable of acting as a replicon or replication intermediate, either independently, or with factors supplied *in trans*. The viral DNA may or may not encode infectious viral particles and furthermore may contain insertions, deletions, substitutions, rearrangements or other modifications. The viral DNA may contain heterologous DNA. In this case, heterologous DNA refers to any non-viral DNA or DNA from a different virus. For example, the heterologous DNA may comprise an expression cassette for a protein or RNA of interest.

Viral replicons suitable for use in the methods and compositions of the invention include those of viruses having a circular DNA genome or replication intermediate, such as: Abutilon mosaic virus (AbMV), African cassava mosaic virus (ACMV), banana streak virus (BSV), bean dwarf mosaic (BDMV), bean golden mosaic virus (BGMV), beet curly top virus (BCTV), beet western yellows virus (BWYV) and other luteoviruses, cassava latent virus (CLV), carnation etched virus (CERV), cauliflower mosaic virus (CaMV), chloris striate mosaic (CSMV),

commelina yellow mottle virus (CoYMV), cucumber mosaic virus (CMV), dahlia mosaic virus (DaMV), digitaria streak virus (DSV), figwort mosaic virus (FMV), hop stunt viroid (HSV), maize streak virus (MSV), mirabilis mosaic virus (MMV), miscanthus streak virus (MiSV), potato stunt tuber virus (PSTV),
5 panicum streak virus (PSV), potato yellow mosaic virus (PYMV), rice tungro bacilliform virus (RTBV), soybean chlorotic mottle virus (SoyCMV), squash leaf curl virus (SqLCV), strawberry vein banding virus (SVBV), sugarcane streak virus (SSV), thistle mottle virus (ThMV), tobacco mosaic virus (TMV), tomato golden mosaic virus (TMGV), tomato mottle virus (TMoV), tobacco ringspot virus
10 (TobRV), tobacco yellow dwarf virus (TobYDV), tomato leaf curl virus (TLCV), tomato yellow leaf curl virus (TYLCV), tomato yellow leaf curl virus-Thailand (TYLCV-t) and wheat dwarf virus (WDV) and derivatives thereof. Preferably the viral replicon is from MSV, WDV, TGMV or TMV.

By "T-DNA" is meant the T-DNA of an *Agrobacterium tumefaciens*
15 Ti plasmid or from an *Agrobacterium rhizogenes* Ri plasmid, or a derivative thereof. The T-DNA may comprise an entire T-DNA, but need only comprise the minimal sequences required *in cis* for transfer (i.e., the right and the left T-DNA border sequences). The T-DNAs of the invention have inserted into them, anywhere between the right and left border sequences, a viral DNA flanked by
20 target sites for a site-specific recombinase. The T-DNA may contain deletions, substitutions and/or additional insertions of DNA other than the viral genome and target sites. Preferably the T-DNA contains an expression cassette for the cognate recombinase of the target sites flanking the viral DNA. The sequences encoding factors required *in trans* for transfer of the T-DNA into a plant cell, such as *vir*
25 genes, may be inserted into the T-DNA, or may be present on the same replicon as the T-DNA, or *in trans* on a compatible replicon in the *Agrobacterium* host. Preferably the *trans* acting factors required for T-DNA transfer are present on the same replicon as the T-DNA.

By "*Agrobacterium* replicon" is meant any replicon (e.g., plasmid
30 or other vector) that is capable of being stably maintained in an *Agrobacterium* host. Such replicons include the *Agrobacterium* chromosome, *Agrobacterium*

plasmids, cosmids, phagemids, etc., derivatives thereof, and any other vector capable of stable replication in *Agrobacterium*. For example, a binary vector suitable for both *Agrobacterium*-mediated transfer and for facile recombinant manipulations and replication in other organisms is useful in the methods and
5 compositions of the invention. Preferably the replicon is a Ti or Ri plasmid or a derivative thereof.

In the compositions and methods of the invention, the viral replicon is flanked by directly repeated target sites for a site-specific recombinase. By "flanked by" is meant that the target sites may be directly contiguous with the viral
10 DNA or there may be one or more intervening sequences present between one or both ends of the viral DNA and the target sites. Intervening sequences of particular interest would include linkers, adapters, selectable markers and/or other sites which aid in vector construction or analysis and expression cassette for a gene of interest. Target sites for site-specific recombinases are known to those skilled in
15 the art and are discussed in co-pending provisional application 60/065,613. Examples of target sites include, but are not limited to FRT, FRT1, FRT5, FRT6, FRT7, other FRT mutants, loxP, loxP mutants, and the like. The loxP and FRT sites are shown in Figure 4.

By "directly repeated" is meant that the target sites that flank the
20 viral DNA are arranged in the same orientation, so that recombination between these sites results in excision, rather than inversion, of the viral DNA.

By "site-specific recombinase" is meant any enzyme capable of being functionally expressed in plants, that catalyzes conservative site-specific recombination between its corresponding target sites. For reviews of site-specific
25 recombinases, see Sauer (1994) *Current Opinion in Biotechnology* 5:521-527; Sadowski (1993) *FASEB* 7:760-767; the contents of which are incorporated herein by reference. Methods of using site-specific recombination systems to excise DNA fragments from chromosomal or extrachromosomal plant DNA are known to those skilled in the art. The bacteriophage P1 *loxP*-Cre and the *Saccharomyces* 2 μ
30 plasmid FRT/FLP site-specific recombinations systems have been extensively studied. For example, Russell *et al.* (1992, *Mol. Gen. Genet.* 234:49-59) describe

the excision of selectable markers from tobacco and *Arabidopsis* genomes using the *loxP*-Cre site-specific recombination system.

It is furthermore recognized that the recombinase, which is used in the invention, will depend upon the target sites of the transformed organism and the targeting cassette. That is, if *FRT* sites are utilized, the FLP recombinase will be needed. In the same manner, where *lox* sites are utilized, the Cre recombinase is required. If the non-identical target sites comprise both a *FRT* and a *lox* site, both the FLP and Cre recombinase or a chimeric recombinase having both Cre and FLP functions, as described in the concurrently filed provisional application entitled "A NOVEL METHOD FOR THE INTEGRATION OF FOREIGN DNA INTO EUKARYOTIC GENOMES," will be required in the plant cell.

of T-DNA to a plant cell, wherein the T-DNA contains a viral replicon flanked by directly repeated target sites for a site-specific recombinase and optionally a DNA of interest linked to the viral replicon. The DNA of interest may also contain a non-identical target site for the recombinase. An expression cassette for the site-specific recombinase is present on the T-DNA or the plant genome, or is transiently introduced into the plant cell. Expression of the site-specific recombinase in the plant cell results in excision of the viral replicon and the associated DNA of interest. The viral replicon and DNA of interest are then replicated to high copy number in the host plant cell.

The compositions of the invention comprise nucleic acids, such as T-DNAs containing a viral DNA flanked by directly repeated target sites for a site-specific The site-specific recombinase may be a naturally occurring recombinase or an active fragment derivative thereof. Site-specific recombinases useful in the methods and compositions of the invention, include recombinases from the integrase and resolvase families, derivatives thereof, and any other naturally occurring or recombinantly produced enzyme or derivative thereof, that catalyzes conservative site-specific recombination between specified DNA sites. The integrase family of recombinases has over thirty members and includes FLP, Cre, Int and R. The resolvase family includes $\gamma\delta$ resolvase. Recombinant enzymes that catalyze site-specific conservative recombination include moFLP, described in co-

-12-

pending application 08/972,258, and moCre, described in the concurrently filed provisional application entitled, "An Artificial Site-Specific DNA Recombinase With Specificities to Multiple Recombination Target Sites"; the contents of which are incorporated herein by reference. Preferably, the recombinase is one that does not require cofactors or a supercoiled substrate. Such recombinases include Cre, FLP moflp, and moCre.

moflp is derived from the *Saccharomyces* 2 μ plasmid FLP recombinase, but is encoded by a nucleic acid sequence utilizing maize-preferred codons. While the *moflp* nucleic acid sequence includes preferred codons for expression of amino acids in maize, it is understood that a useful sequence may contain codons occurring in maize with less than the highest reported maize codon frequencies.

The site-specific recombinases and sequences encoding them that are used in the methods and compositions of the invention may be variants of naturally occurring recombinases and the genes encoding them. The term "conservatively modified variants" applies to both amino acid and nucleic acid sequences. With respect to particular nucleic acid sequences, conservatively modified variants refers to those nucleic acids which encode identical or conservatively modified variants of the amino acid sequences. Because of the degeneracy of the genetic code, a large number of functionally identical nucleic acids encode any given protein. For instance, the codons GCA, GCC, GCG and GCU all encode the amino acid alanine. Thus, at every position where an alanine is specified by a codon, the codon can be altered to any of the corresponding codons described without altering the encoded polypeptide. Such nucleic acid variations are "silent variations" and represent one species of conservatively modified variation. One of ordinary skill will recognize that each codon in a nucleic acid (except AUG, which is ordinarily the only codon for methionine) can be modified to yield a functionally identical molecule.

As to amino acid sequences, one of skill will recognize that individual substitutions, deletions or additions to a nucleic acid, peptide, polypeptide, or protein sequence which alters, adds or deletes a single amino acid

or a small percentage of amino acids in the encoded sequence is a "conservatively modified variant" where the alteration results in the substitution of an amino acid with a chemically similar amino acid. Thus, any number of amino acid residues selected from the group of integers consisting of from 1 to 15 can be so altered.

- 5 Thus, for example, 1, 2, 3, 4, 5, 7, or 10 alterations can be made. Conservatively modified variants typically provide similar biological activity as the unmodified polypeptide sequence from which they are derived. For example, substrate specificity, enzyme activity, or ligand/receptor binding is generally at least 30%, 40%, 50%, 60%, 70%, 80%, or 90% of the native protein for its native substrate.
- 10 Conservative substitution tables providing functionally similar amino acids are well known in the art.

The following six groups each contain amino acids that are conservative substitutions for one another:

- 1) Alanine (A), Serine (S), Threonine (T);
- 15 2) Aspartic acid (D), Glutamic acid (E);
- 3) Asparagine (N), Glutamine (Q);
- 4) Arginine (R), Lysine (K);
- 5) Isoleucine (I), Leucine (L), Methionine (M), Valine (V); and
- 6) Phenylalanine (F), Tyrosine (Y), Tryptophan (W).

- 20 See Creighton (1984) *Proteins*, W.H. Freeman and Company.

- When the nucleic acid is prepared or altered synthetically, advantage can be taken of known codon preferences of the intended host where the nucleic acid is to be expressed. For example, although nucleic acid sequences of the present invention may be expressed in both monocotyledonous and
- 25 dicotyledonous plant species, sequences can be modified to account for the specific codon preferences and GC content preferences of monocotyledons or dicotyledons as these preferences have been shown to differ (Murray *et al.* (1989) *Nucl. Acids Res.* 17:477-498; and Campbell *et al.* (1990) *Plant Physiol.* 92:1). Thus, the maize preferred codon for a particular amino acid may be derived from known gene
- 30 sequences from maize. Maize codon usage for 28 genes from maize plants are listed in Table 4 of Murray *et al.*, *supra*.

The FLP recombinase gene from yeast (*Saccharomyces cerevisiae*) is commercially available in plasmid pOG44 from Stratagene Cloning Systems (11011 North Torrey Pines Road, La Jolla, CA 92037). Similarly, the sequences of many other site specific recombinases and their cognate target sites are publicly or commercially available.

Genes encoding site-specific recombinases, viral DNA, T-DNA and target sites can be made using (a) standard recombinant methods, (b) synthetic techniques, or combinations thereof. Use of cloning vectors, expression vectors, adapters, and linkers is well known in the art and can be found in such references as Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, 2nd ed. (Cold Spring Harbor, New York, 1989). A variety of strategies are available for ligating fragments of DNA, the choice of which depends on the nature of the termini of the DNA fragments and which choices can be readily made by those of skill in the art.. For a description of various nucleic acids see, for example, Stratagene Cloning Systems, Catalogs 1995, 1996, 1997 (La Jolla, CA); and, Amersham Life Sciences, Inc, Catalog =97 (Arlington Heights, IL). Genes encoding FLP can be obtained, for example, by synthesizing the genes with mutually priming long oligonucleotides. See, for example, Ausubel *et al.* (eds.), *Current Protocols In Molecular Biology*, pages 8.2.8 to 8.2.13, Wiley Interscience (1990). Also, see Wosniak *et al.* (1987) *Gene* 60:115. Moreover, current techniques using the polymerase chain reaction provide the ability to synthesize genes as large as 1.8 kilobases in length (Adang *et al.* (1993) *Plant Mol. Biol.* 21:1131; Bombat *et al.* (1993) *PCR Methods and Applications* 2:266).

Rather than use full length recombinases, functional fragments of site-specific recombinases may be used in the methods and compositions of the invention. Functional fragments of site-specific recombinases can be identified using a variety of techniques such as restriction analysis, Southern analysis, primer extension analysis, and DNA sequence analysis. Primer extension analysis or S1 nuclease protection analysis, for example, can be used to localize the putative start site of transcription of the cloned gene. Ausubel at pages 4.8.1 to 4.8.5; Walmsley *et al.*, *Quantitative and Qualitative Analysis of Exogenous Gene Expression by the*

S1 Nuclease Protection Assay, in Methods In Molecular Biology, Vol. 7: Gene Transfer and Expression Protocols, Murray (ed.), pages 271-281 (Humana Press, Inc. 1991). For examples, functional fragments of the FLP protein may be identified by their ability, upon introduction to cells containing appropriate FRT
5 substrates, to catalyze site-specific recombination (for example, excision of an FRT-flanked sequence which upon removal will activate an assayable marker gene.

The general approach of such functional analysis involves subcloning DNA fragments of a genomic clone, cDNA clone or synthesized gene
10 sequence into an expression vector, introducing the expression vector into a heterologous host, and screening to detect the product of recombination (*i.e.* using restriction analysis to verify the product of recombination at the nucleic acid level, or relying on an assay system for recombination as described above). Methods for generating fragments of a CDNA or genomic clone are well known. Variants of an
15 isolated DNA encoding a site-specific recombinase can be produced by deleting, adding and/or substituting nucleotides. Such variants can be obtained, for example, by oligonucleotide-directed mutagenesis, linker-scanning mutagenesis, mutagenesis using the polymerase chain reaction, and the like. See, for example, Ausubel, pages 8.0.3 - 8.5.9. Also, see generally, McPherson (ed.), Directed
20 Mutagenesis: A Practical approach, (IRL Press, 1991).

The nucleic acid sequence encoding the site-specific recombinase is operably linked to a promoter that drives expression in a plant. As used herein "operably linked" includes reference to a functional linkage between a promoter and a second sequence, wherein the promoter sequence initiates and mediates
25 transcription of the DNA sequence corresponding to the second sequence. Generally, operably linked means that the nucleic acid sequences being linked are contiguous and, where necessary to join two protein coding regions, contiguous and in the same reading frame.

As used herein "promoter" includes reference to a region of DNA
30 upstream from the start of transcription and involved in recognition and binding of RNA polymerase and other proteins to initiate transcription. A "plant promoter" is

a promoter capable of initiating transcription in plant cells. Exemplary plant promoters include, but are not limited to, those that are obtained from plants, plant viruses, and bacteria genes that are expressed in plant cells such as those of *Agrobacterium* or *Rhizobium*. Both heterologous and non-heterologous (i.e., endogenous) promoters can be employed to direct expression of a sequence encoding a site-specific recombinase. The promoter may be constitutive, inducible or tissue specific.

Many different constitutive promoters can be utilized in the instant invention. Exemplary constitutive promoters include the promoters from plant viruses such as the 35S promoter from CaMV (Odell *et al.* (1985) *Nature* 313:810-812) and the promoters from such gene as rice actin (McElroy *et al.* (1990) *Plant Cell* 2:163-171); ubiquitin (Christensen *et al.* (1989) *Plant Mol. Biol.* 12:619-632 and Christensen *et al.* (1992) *Plant Mol. Biol.* 18:675-689); pEMU (Last *et al.* (1991) *Theor. Appl. Genet.* 81:581-588); MAS (Velten *et al.* (1984) *EMBO J.* 3:2723-2730); maize H3 histone (Lepetit *et al.* (1992) *Mol. Gen. Genet.* 231: 276-285 and Atanassova *et al.* (1992) *Plant Journal* 2(3):291-300); the 1'- or 2'- promoter derived from T-DNA of *Agrobacterium tumefaciens*, the ubiquitin 1 promoter, the Smas promoter, the cinnamyl alcohol dehydrogenase promoter (U.S. Patent No. 5,683,439), the *Nos* promoter, the Pemu promoter, the rubisco promoter, the GRP1-8 promoter, and other transcription initiation regions from various plant genes known to those of skill. The ALS promoter, a XbaI/NcoI fragment 5-prime to the *Brassica napus* ALS3 structural gene (or a nucleotide sequence that has substantial sequence similarity to said XbaI/NcoI fragment), represents a particularly useful constitutive promoter for dicots. (See co-pending Pioneer Hi-Bred International US Patent Application 08/409,297).

A variety of inducible promoters can be used in the instant invention. See Ward *et al.* (1993) *Plant Mol. Biol.* 22:361-366. Exemplary inducible promoters include that from the ACE1 system which responds to copper (Mett *et al.* (1993) *PNAS* 90:4567-4571); In2 gene from maize which responds to benzenesulfonamide herbicide safeners (Hershey *et al.* (1991) *Mol. Gen. Genetics* 227:229-237 and Gatz *et al.* (1994) *Mol. Gen. Genetics* 243:32-38); the Adh1

promoter which is inducible by hypoxia or cold stress, the Hsp70 promoter which is inducible by heat stress, and the PPDk promoter which is inducible by light; or Tet repressor from Tn10 (Gatz *et al.* (1991) *Mol. Gen. Genet.* 227:229-237. A particularly preferred inducible promoter is a promoter that responds to an inducing agent to which plants do not normally respond. An exemplary inducible promoter is the inducible promoter from a steroid hormone gene the transcriptional activity of which is induced by a glucocorticosteroid hormone (Schena *et al.* (1991) *Proc. Natl. Acad. Sci. U.S.A.* 88:10421).

Examples of promoters under developmental control include promoters that initiate transcription only, or preferentially, in certain tissues, such as leaves, roots, fruit, seeds, or flowers. The operation of a promoter may also vary depending on its location in the genome. Thus, an inducible promoter may become fully or partially constitutive in certain locations.

The site-specific recombinase must be expressed in the plant cell in order for mobilization of the viral DNA from the T-DNA. Accordingly, the recombinase protein must be expressed so that it is present in the cell at an effective concentration sometime between when the T-DNA is transferred to the plant cell and before non-integrated T-DNA is lost from the plant cells.

Accordingly, the expression cassette encoding the site-specific recombinase may be supplied on the T-DNA *in cis* to the viral DNA; *in trans* on a plant chromosome or extrachromosomal replicon; or may be transferred to the plant near to the time of agro-mediated transformation. The site-specific recombinase gene may be constitutively or transiently expressed, or may be inducible. If cofactors are required for a particular site-specific recombination system, they may be encoded on the T-DNA of the invention or in the plant host chromosomal or extrachromosomal DNA.

By "target site for a site-specific recombinase" is meant a DNA sequence that is recognized by a particular site-specific recombinase. A variety of target sites are known to those skilled in the art and may be used in the methods and compositions of the invention. The site may have the sequence of the cognate site for a given recombinase, or may be modified, so long as it is capable of acting

as a target site. The site may contain the minimal sequences necessary for recombination, or it may contain additional sequences that enhance recombination. Examples of target sites for use in the invention are known in the art and include FRT and *loxP* sites (See, for example, Schlake and Bode (1994) *Biochemistry* 33:12746-12751; Huang *et al.* (1991) *Nucleic Acids Research* 19:443-448; Paul D. Sadowski (1995) *In Progress in Nucleic Acid Research and Molecular Biology* vol. 51, pp. 53-91; Michael M. Cox (1989) *In Mobile DNA*, Berg and Howe (eds) American Society of Microbiology, Washington D.C., pp. 116-670; Dixon *et al.* (1995) 18:449-458; Umlauf and Cox (1988) *The EMBO Journal* 7:1845-1852; Buchholz *et al.* (1996) *Nucleic Acids Research* 24:3118-3119; Kilby *et al.* (1993) *Trends Genet.* 9:413-421; Rossant and Geagy (1995) *Nat. Med.* 1: 592-594; *Lox* Albert *et al.* (1995) *The Plant J.* 7:649-659; Bayley *et al.* (1992) *Plant Mol. Biol.* 18:353-361; Odell *et al.* (1990) *Mol. Gen. Genet.* 223:369-378; and Dale and Ow (1991) *Proc. Natl. Acad. Sci. USA* 88:10558-105620; Qui *et al.* (1994) *Proc. Natl. Acad. Sci. USA* 91:1706-1710; Stuurman *et al.* (1996) *Plant Mol. Biol.* 32:901-913; Hartley *et al.* (1980) *Nature* 286: 860-864; Sauer (1994) *Current Opinion in Biotechnology* 5:521-527; and Dale *et al.* (1990) *Gene* 91:79-85; all of which are herein incorporated by reference.)

Each *loxP* and FRT site contains two 13 base pair inverted repeats which flank an 8 base pair spacer. The FRT site contains an additional non-essential 13 base pair repeat. The sequences of the *loxP* and FRT sites are shown in Figure 1. A minimal FRT site comprising two 13 base pair repeats, separated by an 8 base spacer, is:

5'-GAAGTTCCTATTC[TCTAGAAA]GTATAGGAACTTC3'

wherein the nucleotides within the brackets indicate the spacer region. The nucleotides in the spacer region can be replaced with a combination of nucleotides, so long as the two 13-base repeats are separated by eight nucleotides. FLP is a conservative, site-specific recombinase, capable of catalyzing inversion of a nucleic acid sequence positioned between two inversely oriented FRTs; recombination between two molecules each containing a FRT site; and excision between FRT sites. The core region is not symmetrical, and its asymmetry dictates

the directionality of the reaction. Recombination between inverted FRT sites causes inversion of a DNA sequence between them, whereas recombination between directly oriented sites leads to excision of the DNA between them.

5 T-DNAs containing viral DNA flanked by recombinations sites, expression cassettes for site-specific recombinases and vectors carrying these sequences may be constructed using standard molecular biology techniques. See, for example, Sambrook *et al.* (eds.) *Molecular Cloning: A Laboratory Manual*, Second Edition, (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY 1989).

10 Typical vectors useful in the methods and compositions of the invention are well known in the art and include vectors derived from the tumor-inducing (Ti) plasmid of *Agrobacterium tumefaciens* described by Rogers *et al.* (1987) *Meth. in Enzymol.*, 153:253-277. These vectors are plant integrating vectors in that on transformation, the vectors integrate a portion of vector DNA
15 into the genome of the host plant. Exemplary *A. tumefaciens* vectors useful herein are plasmids pKYLX6 and pKYLX7 of Schardl *et al.*, *Gene*, 61:1-11 (1987) and Berger *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 86:8402-8406 (1989). Another useful vector herein is plasmid pBI101.2 that is available from Clontech Laboratories, Inc. (Palo Alto, CA).

20 Techniques for transforming a wide variety of higher plant species are well known and described in the technical, scientific, and patent literature. See, for example, Weising *et al.*, *Ann. Rev. Genet.* 22: 421-477 (1988). These methods are useful for transforming a plant cell with a site-specific recombinase expression cassette. This step will be necessary if the cassette is not included in the
25 transferred T-DNA. The expression cassette encoding the site-specific recombinase may be present in the plant genome prior to Agroinfection or may be transformed into the plant around the time of transfer of the T-DNA to the plant cell so that it will be transiently expressed.

For example, the DNA construct may be introduced directly into the genomic
30 DNA of the plant cell using techniques such as electroporation, PEG poration, particle bombardment, silicon fiber delivery, microinjection of plant cell

protoplasts or embryogenic callus, or *Agrobacterium*-mediated transformation (Hiei *et al.* (1994) *Plant J.* 6:271-282).

The introduction of DNA constructs using polyethylene glycol precipitation is described in Paszkowski *et al.*, *Embo J.* 3: 2717-2722 (1984).

- 5 Electroporation techniques are described in Fromm *et al.*, *Proc. Natl. Acad. Sci.* 82: 5824 (1985). Ballistic transformation techniques are described in Klein *et al.*, *Nature* 327: 70-73 (1987).

DNA can also be introduced into plants by direct DNA transfer into pollen as described by Zhou *et al.*, *Methods in Enzymology*, 101:433 (1983); D.

- 10 Hess, *Intern Rev. Cytol.*, 107:367 (1987); Luo *et al.*, *Plane Mol. Biol. Reporter*, 6:165 (1988). Expression of polypeptide coding genes can be obtained by injection of the DNA into reproductive organs of a plant as described by Pena *et al.*, *Nature*, 325.:274 (1987). DNA can also be injected directly into the cells of immature embryos and the rehydration of desiccated embryos as described by Neuhaus *et al.*,
15 *Theor. Appl. Genet.*, 75:30 (1987); and Benbrook *et al.*, in *Proceedings Bio Expo* 1986, Butterworth, Stoneham, Mass., pp. 27-54 (1986). A variety of plant viruses that can be employed as vectors are known in the art and include cauliflower mosaic virus (CaMV), geminivirus, brome mosaic virus, and tobacco mosaic virus.

- Plants cells stably transformed with a site-specific recombinase
20 expression cassette can be regenerated, e.g., from single cells, callus tissue or leaf discs according to standard plant tissue culture techniques. It is well known in the art that various cells, tissues, and organs from almost any plant can be successfully cultured to regenerate an entire plant. Plant regeneration from cultured protoplasts is described in Evans *et al.*, *Protoplasts Isolation and Culture, Handbook of Plant*
25 *Cell Culture*, Macmillilan Publishing Company, New York, pp. 124-176 (1983); and Binding, *Regeneration of Plants, Plant Protoplasts*, CRC Press, Boca Raton, pp. 21-73 (1985).

- The regeneration of plants containing the recombinase gene introduced by *Agrobacterium* from leaf explants can be achieved as described by
30 Horsch *et al.*, *Science*, 227:1229-1231 (1985). In this procedure, transformants are grown in the presence of a selection agent and in a medium that induces the

-21-

regeneration of shoots in the plant species being transformed as described by Fraley *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 80:4803 (1983). This procedure typically produces shoots within two to four weeks and these transformant shoots are then transferred to an appropriate root-inducing medium containing the selective agent and an antibiotic to prevent bacterial growth. Transgenic plants of the present invention may be fertile or sterile.

Regeneration can also be obtained from plant callus, explants, organs, or parts thereof. Such regeneration techniques are described generally in Klee *et al.*, *Ann. Rev. of Plant Phys.* 38: 467-486 (1987). The regeneration of plants from either single plant protoplasts or various explants is well known in the art. See, for example, *Methods for Plant Molecular Biology*, A. Weissbach and H. Weissbach, eds., Academic Press, Inc., San Diego, Calif. (1988). This regeneration and growth process includes the steps of selection of transformant cells and shoots, rooting the transformant shoots and growth of the plantlets in soil. For maize cell culture and regeneration see generally, *The Maize Handbook*, Freeling and Walbot, Eds., Springer, New York (1994); *Corn and Corn Improvement*, 3rd edition, Sprague and Dudley Eds., American Society of Agronomy, Madison, Wisconsin (1988).

One of skill will recognize that after the site-specific recombinase expression cassette is stably incorporated in transgenic plants and confirmed to be operable, it can be introduced into other plants by sexual crossing. Any of a number of standard breeding techniques can be used, depending upon the species to be crossed.

The methods and compositions of the invention are useful to mobilize viral DNA from T-DNAs transferred by Agrobacterium to any plant host. As used herein, the term "plant" includes reference to whole plants, plant organs (e.g., leaves, stems, roots, etc.), seeds and plant cells and progeny of same. Plant cell, as used herein includes, without limitation, seeds suspension cultures, embryos, meristematic regions, callus tissue, leaves, roots, shoots, gametophytes, sporophytes, pollen, and microspores. The class of plants which can be used in the methods of the invention is generally as broad as the class of

higher plants amenable to *Agrobacterium* transformation techniques, including both monocotyledonous and dicotyledonous plants. A particularly preferred monocot is maize. Other monocots of particular interest include wheat, rice, barley, sorghum and rye. Dicots of particular interest include soybean, Brassica, sunflower, alfalfa, and safflower.

The T-DNA containing the viral DNA flanked by sites for a site-specific recombinase is transferred to a plant cell by agroinfection. *Agrobacterium tumefaciens*-mediated transformation techniques are well described in the scientific literature. See, for example Horsch *et al.*, *Science* 233: 496-498 (1984), Fraley *et al.*, *Proc. Natl. Acad. Sci.* 80: 4803 (1983) and Kado, (1991), *Crit. Rev. Plant Sci.* 10:1. Descriptions of the *Agrobacterium* vector systems and methods for *Agrobacterium*-mediated gene transfer are provided in Gruber *et al.*, *supra*; Miki, *et al.*, *supra*; and Moloney *et al.* (1989), *Plant Cell Reports* 8:238. Although *Agrobacterium* is useful primarily in dicots, certain monocots can be transformed by *Agrobacterium*. For instance, *Agrobacterium* transformation of maize is described in U.S. Patent No. 5,550,318. Other methods of agroinfection include *Agrobacterium rhizogenes*-mediated transformation (see, e.g., Lichtenstein and Fuller In: Genetic Engineering, vol. 6, PWJ Rigby, Ed., London, Academic Press, 1987; and Lichtenstein, C. P., and Draper, J., In: DNA Cloning, Vol. II, D. M. Glover, Ed., Oxford, IRI Press, 1985), Application PCT/US87/02512 (WO 88/02405 published Apr. 7, 1988) describes the use of *A. rhizogenes* strain A4 and its Ri plasmid along with *A. tumefaciens* vectors pARC8 or pARC16.

Optimized methods and vectors for *Agrobacterium*-mediated transformation of plants in the family Graminae, such as rice and maize have been described by Heath *et al.* (1997) *Mol. Plant-Microbe Interact.* 10:221-227; Hiei *et al.* (1994) *Plant J.* 6:271-282 and Ishida *et al.* (1996) *Nat. Biotech.* 14:745-750, the contents of which are incorporated herein by reference. The efficiency of maize transformation is affected by a variety of factors including the types and stages of tissue infected, the concentration of *Agrobacterium*, the tissue culture media, the Ti vectors and the maize genotype. Super binary vectors carrying the vir genes of *Agrobacterium* strains A281 and A348 are useful for high efficiency

transformation of monocots. However, even without the use of high efficiency vectors, it has been demonstrated that T-DNA is transferred to maize at an efficiency that results in systemic infection by viruses introduced by agroinfection, although tumors are not formed (Grimsley *et al.* (1989) *Mol. Gen. Genet.* 217:309-
5 316, the contents of which are incorporated herein by reference). This is because integration of the T-DNA containing the viral genome is not required for viral multiplication, since the excised viral genome acts as an independent replicon.

Another useful basic transformation protocol involves a combination of wounding by particle bombardment, followed by use of
10 *Agrobacterium* for DNA delivery, as described by Bidney *et al.* (1992) *Plant Mol. Biol.* 18:301-313. In general, the intact meristem transformation method involves imbibing seed for 24 hours in the dark, removing the cotyledons and root radical, followed by culturing of the meristem explants. Twenty-four hours later, the primary leaves are removed to expose the apical meristem. The explants are
15 placed apical dome side up and bombarded, *e.g.*, twice with particles, followed by co-cultivation with *Agrobacterium*. To start the co-cultivation for intact meristems, *Agrobacterium* is placed on the meristem. After about a 3-day co-cultivation period the meristems are transferred to culture medium with selection agents.

20 The split meristem method involves imbibing seed, breaking of the cotyledons to produce a clean fracture at the plane of the embryonic axis, excising the root tip and then bisecting the explants longitudinally between the primordial leaves. The two halves are placed cut surface up on the medium then bombarded twice with particles, followed by co-cultivation with *Agrobacterium*. For split
25 meristems, after bombardment, the meristems are placed in an *Agrobacterium* suspension for 30 minutes. They are then removed from the suspension onto solid culture medium for three day co-cultivation. After this period, the meristems are transferred to fresh medium with cefotaxime (plus selection agents).

The excision of viral DNA from the T-DNA may be determined by
30 a variety of means including but not limited to observation of viral lesions or other signs of viral infection on the plant, by antibody-based detection of viral proteins,

by amplification of PCR products specific to a circular genome, or by Northern analysis for viral RNA. In addition, assays for FLP recombinase activity are known and generally measure the overall activity of the enzyme on DNA substrates containing FRT sites. In this manner, a frequency of excision of the sequence can be determined. Excision of DNA from a linear molecule or intermolecular recombination frequency induced by the enzyme may be assayed, as described, for example, in Babineau *et al.* (1985) *JBC* 260:12313; Meyer-Leon *et al.* (1987) *NA Res* 15:6469; and Gronostajski *et al.* (1985) *JBC* 260:12328.

The following examples are offered by way of illustration not by way of limitation.

EXPERIMENTAL

Example 1

Construction of T-DNA Vectors Containing Viral DNA Flanked By Site-Specific Recombination Sites

DNA fragments containing viral DNA flanked by FRT or *loxP* site-specific recombination sites are constructed either by synthesizing, annealing and ligating complementary oligonucleotides or by creating primers for PCR amplification of a DNA product containing the FRT or *loxP* site and restriction sites useful for cloning into T-DNA near the 5' end of the PCR product.

For example, long PCR primers may be designed wherein the 3' end of the primer hybridizes to the 5' end of the viral genome of interest and the 5' end of the primer further contains a *loxP* site and useful cloning sites (see Figure 5). The resulting PCR product is digested with the appropriate restriction enzyme and inserted into the T-DNA of an *Agrobacterium* binary vector as described by Bevan, M. (1984) *Nucl. Acids Res.* 12:8711-8721; herein incorporated by reference. An expression cassette for Cre is also inserted into the T-DNA. The resulting constructs are transformed into *E. coli*, screened, and then transferred to *Agrobacterium*.

Example 2

Transfer of T-DNA to Dicotyledonous Plant Cells

5 *Nicotiana tabacum* BY-2 cells are propagated in Murashige and
Skoog medium (Gibco BRL) containing 3% sucrose, 1 Fg/ml thiamine, 0.2 Fg/ml
2,4-D, and 370 Fg/ml KH_2PO_4 . *Zea mays* Black Mexican Sweet (BMS) cells are
propagated in Murashige and Skoog medium containing 2% sucrose, 2 Fg/ml 2,4-
D, 0.2 mg/ml myoinositol, 0.13 mg/ml L-asparagine, 0.13 Fg/ml nicotinic acid,
10 and 0.25 Fg/ml each of thiamine, pyridoxine, and pantothenic acid. The cultures
are shaken at 140 rpm at 25EC in continuous light.

To infect plant cells, virulence (*vir*) gene activity is induced by
treatment with acetosyringone in *Agrobacterium* carrying the vector constructed in
Example 1. *Agrobacterium* cells are grown to a density of 2×10^9 cells per ml (A
15 = 100, using a Klett-Summerson spectrophotometer, red filter) in AB-sucrose
medium. The cells are centrifuged at 10,000g, suspended at a concentration of $1 \times$
 10^9 cells per ml ($A = 50$) in induction medium (AB salts, 0.5% glucose, 2mM
sodium phosphate, 50 Mm Mes, Ph 5.6, 50FM acetosyringone), and incubated with
gentle shaking at 25EC for 14 to 18 hr. After washing the bacterial cells in plant
20 culture medium, plant cells are inoculated with induced *Agrobacterium* (-20
bacterial cells per plant cell, except where noted otherwise) and cocultivated at
25EC with shaking at 140 rpm for various periods of time. Most of the bacteria is
washed off by centrifugation of the cocultivation mixture at 300 rpm (model GLC-
2 clinical centrifuge; Beckman Sorvall, Newtown, CT) for 2 min. The plant cell
25 pellet is suspended and washed once more in plant culture medium and then
resuspended in culture containing either 100 Fg/ml timentin or 200 Fg/mL
cefotaxime. Mobilization of viral genomes from the T-DNA is monitored by
quantitating infectious viral particles.

Example 3

Transfer of T-DNA to Monocotyledonous Plants

Prior to inoculation, strains of *Agrobacterium* carrying the vector
5 constructed in Example 1 are streaked out on YEB (Grimsley *et al.* (1986) *Proc. Natl. Acad. Sci. USA*, 83:3282-3286) plates solidified with 1.5% agar and supplemented with 100Fg/ml rifampicin and 25Fg/ml kanamycin and allowed to grow for 48h. A single colony is used to inoculate 10 ml of liquid YEB medium in a 100ml Erlenmeyer flask supplemented with antibiotics as previously. Growth is
10 continued with shaking at 200 r.p.m. for 24h, then 500Fl of this culture is used to inoculate a similar flask and growth continued for a further 20h. This procedure yields a final density of viable *Agrobacterium* cells in the region of 10^9 /ml (estimated by plating). The cells are then harvested by centrifugation and resuspended in an equal volume of 10mM $MgSO_4$ without antibiotics; such a
15 suspension is subsequently referred to as undiluted or 10^0 dilution; for experiments involving a dilution series 10mM $MgSO_4$ was also used as the diluent.

Maize seeds for 10-day old plants are sown in pots in a phytotron in a 12 hour light/dark cycle at 25EC in a light intensity of about 10000 lux (Sylvania 215W fluorescent lamps type F96T12/CW/VHO) then moved to the BL3
20 containment laboratory immediately prior to inoculation; subsequent growth conditions have been described (Grimsley *et al.* (1987) *Nature* 325:177-179). Three-day old seedlings are prepared by (i) sterilization by stirring for 20min in 0.7% calcium hypochlorite solution, (ii) washing three times (stirring for 20min each time) in sterile distilled water (iii) preparing 9cm diameter presterilized Petri
25 dishes with 3 sheets of sterile 8.5cm diameter Macherey-Nagel (Germany) filter paper in the bottom and *ca.* 10ml of sterile water per dish, (iv) putting *ca.* 20 seeds into each geranium dish, and (v) incubating in the dark at 28EC for 3 days, or until the distance between the scutellar node and the apical tip of the coleoptiles is 1-2
cm.

30 For inoculation of plants, a 50Fl or a 100Fl Hamilton syringe fitted with a 0.4mm diameter disposable needle is loaded with the bacterial suspension

-27-

avoiding trapped air bubbles. Between inoculations with different bacterial strains the needle is discarded and the syringe flushed out 3 times with 100% ethanol and 3 times with sterile distilled water. 10-day old plants are inoculated by (i) abrasion of an upper leaf, applying 20Fl of suspension, and rubbing in with carborundum powder until the leaf appears wet all over, (ii) injection of 10Fl of bacterial suspension into the central part of the plant either just above the first leaf blade, or 1cm below the first leaf blade, or at the base of the plant, in the meristematic region where adventitious roots later begin to appear. Three-day old seedlings are injected with 10Fl of bacterial suspension in different ways by (i) pushing the needle down through the apical tip of the coleoptile to the coleoptilar node, (ii) injecting 2mm below the apical tip of the coleoptile, (iii) 2 mm above the coleoptilar node, (iv) at the coleoptilar node, (iv) 2mm below the coleoptilar node, (v) at the scutellar node, and by pushing the needle up through the primary root to a region close to the scutellar node. Ten Fl is used as a standard inoculum of bacterial suspension, but only 1-2Fl routinely remains in the inoculation site, the rest is forced out, usually coming out from the point of entry of the inoculating needle. Following inoculation seedlings are planted immediately in damp soil, incubated as before (Grimsley *et al.*(1987) *Nature* 325:177-179), and observed daily for the appearance of symptoms of viral infection, characterized by the appearance of yellow spots and/or stripes at the base of new leaves.

All publications and patent applications mentioned in the specification are indicative of the level of those skilled in the art to which this invention pertains. All publications and patent applications are herein incorporated by reference to the same extent as if each individual publication or patent application was specifically and individually indicated to be incorporated by reference.

Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be obvious that certain changes and modifications may be practiced within the scope of the appended claims.

THAT WHICH IS CLAIMED:

1. A method for mobilizing a viral replicon from a T-DNA,
comprising:
 - a) providing an *Agrobacterium* replicon having a T-DNA, wherein said T-DNA contains a viral replicon flanked by directly repeated
5 target sites for a site-specific recombinase; and
 - b) infecting a cell of a plant with an *Agrobacterium*
carrying said *Agrobacterium* replicon under conditions that allow transfer of said
T-DNA and expression of said recombinase in said cell;
wherein said cell, said T-DNA, or said viral replicon
10 contains a nucleotide sequence encoding said recombinase or an active fragment or
variant thereof, and said nucleotide sequence is operably linked to a promoter that
drives expression in said cell.
2. A method for providing to a plant cell a plurality of copies
15 of a DNA sequence of interest, comprising:
 - a) providing an *Agrobacterium* replicon having a T-DNA, wherein said T-DNA contains in a 5' to 3' or 3' to 5' orientation, a first target
site for a site-specific recombinase, a viral replicon, said DNA sequence of interest,
and a second target site for said recombinase in direct repeat with said first target
20 site, wherein said first and second target sites are identical; and
 - b) infecting a cell of a plant with an *Agrobacterium*
carrying said *Agrobacterium* replicon under conditions that allow transfer of said
T-DNA and expression of said recombinase in said cell;
wherein said cell, said T-DNA, or said viral replicon
25 contains a nucleotide sequence encoding said recombinase or an active fragment or
variant thereof, and said nucleotide sequence is operably linked to a promoter that
drives expression in said cell.
3. A method for providing to a plant cell a plurality of copies

of a DNA sequence of interest flanked by nonidentical target sites for a site-specific recombinase, comprising:

- a) providing an *Agrobacterium* replicon having a T-DNA, wherein said T-DNA contains in a 5' to 3' or 3' to 5' orientation, a first target site for said recombinase, a viral replicon, a second target site for said recombinase, said DNA sequence of interest, and a third target site for said recombinase, wherein said first and third target sites are directly repeated and identical with respect to each other, and said second target site is nonidentical to said first and third target sites; and
- b) infecting a cell of a plant with an *Agrobacterium* carrying said *Agrobacterium* replicon under conditions that allow transfer of said T-DNA and expression of said recombinase in said cell;
wherein said cell, said T-DNA, or said viral replicon contains a nucleotide sequence encoding said recombinase or an active fragment or variant thereof, and said nucleotide sequence is operably linked to a promoter that drives expression in said cell.

4. The method of any one of claims 1-3, wherein said site-specific recombinase is a member of the integrase family.

20

5. The method of claim 4, wherein said site-specific recombinase is FLP, Cre, Int, SSVI integrase R or an active fragment or variant thereof.

25

6. The method of claim 5, wherein said target sites are selected from the group consisting of FRT and mutant FRT sites.

7. The method of claim 6, wherein said mutant FRT site is FRT5, FRT6 or FRT7.

30

8. The method of claim 6, wherein said recombinase is encoded by moFLP.

-30-

9. The method of claim 5, wherein said target sites are loxP and a mutant loxP.

5 10. The method of claim 9, wherein said recombinase is encoded by moCre.

11. The method of claim 3, wherein said *Agrobacterium* replicon is a Ti plasmid or a derivative thereof.

10

12. The method of claim 3, wherein said *Agrobacterium* replicon is a Ri plasmid or a derivative thereof.

13. The method of claim 3, wherein said T-DNA contains said nucleotide sequence encoding said recombinase or an active fragment or variant thereof.

15

14. The method of claim 3, wherein said viral replicon contains said nucleotide sequence encoding said recombinase or an active fragment or variant thereof.

20

15. The method of claim 3, wherein said plant is a monocot.

16. The method of claim 15, wherein said monocot is a member of the family *Graminae*.

25

17. The method of claim 15, wherein said monocot is maize, wheat, rice, barley, sorghum or rye.

18. The method of claim 3, wherein said viral replicon is a MSV, WDV, TGMV or TMV replicon.

30

-31-

19. The method of claim 3, wherein said plant is a dicot.

20. The method of claim 19, wherein said dicot is soybean, canola, Brassica, sunflower, alfalfa or safflower.

5

21. A T-DNA containing a viral replicon flanked by directly repeated target sites for a site-specific recombinase.

22. A T-DNA containing in a 5' to 3' or 3' to 5' orientation, a first target site for a site-specific recombinase, a viral replicon, said DNA sequence of interest, and a second target site for said recombinase in direct repeat with said first target site, wherein said first and second target sites are identical.

23. A T-DNA containing in a 5' to 3' or 3' to 5' orientation, a first target site for said recombinase, a viral replicon, a second target site for said recombinase, said DNA sequence of interest, and a third target site for said recombinase, wherein said first and third target sites are directly repeated and identical with respect to each other, and said second target site is nonidentical to said first and third target sites.

20

24. The T-DNA of claim 23, wherein said viral replicon contains a nucleotide sequence encoding said recombinase or an active fragment or variant thereof, and said nucleotide sequence is operably linked to a promoter that drives expression in a plant cell.

25

25. The T-DNA of claim 24, wherein said recombinase is encoded by FLP or moFLP.

26. The T-DNA of claim 24, wherein said recombinase is encoded by Cre or moCre.

30

27. The T-DNA of claim 23, wherein said site-specific

-32-

recombinase is a member of the integrase family.

28. The T-DNA of claim 27, wherein said site-specific recombinase is FLP, Cre, Int, R or an active fragment or variant thereof.

5

29. The T-DNA of claim 28, wherein said target sites are selected from the group consisting of FRT and mutant FRT sites.

30. The T-DNA of claim 29, wherein said mutant FRT site is
10 FRT5, FRT6 or FRT7.

31. The T-DNA of claim 28, wherein said target sites are loxP and a mutant loxP.

15 32. The T-DNA of claim 23, further comprising a nucleotide sequence encoding said site-specific recombinase, wherein said nucleotide sequence is operably linked to a promoter that drives expression in a plant cell.

20 33. An Agrobacterium replicon containing the T-DNA of claims 21-32.

34. An Agrobacterium containing the replicon of claim 33.

1/5




FIG. 1.

2/5






FIG. 2.

3/5

FIG. 3.



4/5

FIG. 4.

5/5




FIG. 5.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/24486

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/83 C12N15/84 C12N15/63 C12N5/10 A01H4/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	HAYES, R.J. ET AL.: "Agroinfection of Nicotiana spp. with cloned DNA of tomato golden mosaic virus" THE JOURNAL OF GENERAL VIROLOGY, vol. 69, no. 7, 1988, pages 1487-1496, XP002096355 see the Introduction ---	1-34
A	GB 2 174 995 A (CIBA-GEIGY AG) 19 November 1986 see abstract and page 2, lines 41-43 --- -/--	1-34

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

15 March 1999

Date of mailing of the international search report

20/04/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Alt, G

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/24486

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>SHEN, W.-H. AND HOHN, B.: "Excision of a transposable element from a viral vector introduced into maize plants by agroinfection"</p> <p>THE PLANT JOURNAL, vol. 2, no. 1, 1992, pages 35-42, XP002096356 see the whole document, especially the abstract, lines 4-9</p> <p>---</p>	1-34
A	<p>WO 97 37012 A (COMMONWEALTH SCIENTIFIC AND INDUSTRIAL RESEARCH ORGANISATION) 9 October 1997 see the whole document, especially page 6, lines 9-16</p> <p>---</p>	1-34
A	<p>LYZNIK, L. A. ET AL.: "FLP-mediated recombination of FRT sites in the maize genome"</p> <p>NUCLEIC ACIDS RESEARCH, vol. 24, no. 19, 1996, pages 3784-3789, XP002096357 see the whole document, especially the abstract, lines 3-6</p> <p>---</p>	1-34
A	<p>RUSSELL, S.H. ET AL.: "Directed excision of a transgene from the plant genome"</p> <p>MOL. GENET. GENET., vol. 234, 1992, pages 49--59, XP002096358 cited in the application see the whole document</p> <p>---</p>	1-34
A	<p>WO 96 04393 A (DELTA AND PINE LAND COMPANY) 15 February 1996 see page 16, lines 6-18</p> <p>-----</p>	1-34

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 98/24486

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
GB 2174995	A	19-11-1986	AT 88504 T	15-05-1993
			AU 595527 B	05-04-1990
			AU 5735686 A	20-11-1986
			BR 8602126 A	13-01-1987
			DD 279503 A	06-06-1990
			DE 3688301 D	27-05-1993
			EP 0201904 A	20-11-1986
			JP 61260886 A	19-11-1986
			US 5569597 A	29-10-1996
WO 9737012	A	09-10-1997	AU 2143797 A	22-10-1997
			CA 2250111 A	09-10-1997
WO 9604393	A	15-02-1996	US 5723765 A	03-03-1998
			AU 696668 B	17-09-1998
			AU 3205095 A	04-03-1996
			BR 9508471 A	28-10-1997
			CA 2196410 A	15-02-1996
			EP 0775212 A	28-05-1997
			JP 10503377 T	31-03-1998
			ZA 9506410 A	11-03-1996